SYBR Sa DNAG Sal

	- · · .	and a constant	· · · ·	, 175		
	*, 44 –	· ,¥ I ,,-, , , I ,,, ,∕,		, 9		
	*, 44 –	, 4 1 , -, -, , 1 , , , , , 1				
2	4,	,				
		•				
$ = \sum_{i=1}^{n} \sum_{j=1}^{n} \sum_{i=1}^{n} \sum$						
$[\cdot, \cdot]_{\mathcal{S}_{\mathbf{s}}} := \{\cdot, \cdot\}, I_{\mathcal{S}_{\mathbf{s}}}, I_{\mathcal{S}_{\mathbf{s}}}, \dots, I_{\mathcal{S}_{\mathbf{s}}}\}$						

e og e 🗠 Nee

SYBR®Safe DNA gel stain has been specifically developed for reduced mutagenicity, making it safer than ethidium bromide for staining DNA in agarose or acrylamide gels. SYBR®Safe stain comes either as a concentrate or as a ready-to-use solution that can be used just like an ethidium bromide solution, and the detection sensitivity with SYBR®Safe stain is comparable to that obtained with ethidium bromide. DNA bands stained with SYBR®Safe DNA gel stain can be detected using a standard UV transilluminator, a visible-light transilluminator, or a laser-based scanner. The stain is also suitable for staining RNA in gels. Bound to nucleic acids, SYBR®Safe stain has fluorescence excitation maxima at 280 and 502 nm, and an emission maximum at 530 nm (Figure 1).

	•	•		
. (.)	/ 🛃	/ • • /		
	Sec. Sec. Sec.			
I, <u>, , , , , , , , , , , , , , , , , , </u>) /	.64		
······································	,	4 /		
······································	<u>\</u>			
······································		a second a second second		
J	Sec. Sec. Sec.	. 		
, Lin, (M. , Lin, I., , , (🖉)	Sec. Sec. Sec.	a second a second second		
· · · · · · · · · · · · · · · · · · ·	Sec. Contractor			
	Sec. Sec. Sec.	a second a second se		
· − · · · · · · · · · · · · · · · · · ·				